

MARCH 1981

PPPL-1783


UC-20g

HAMILTONIAN FIELD DESCRIPTION OF
TWO-DIMENSIONAL VORTEX FLUIDS AND
GUIDING CENTER PLASMAS

BY

P. J. MORRISON

PLASMA PHYSICS
LABORATORY



PRINCETON UNIVERSITY
PRINCETON, NEW JERSEY

This work supported by the U.S. Department of Energy
Contract No. DE-AC02-76-CH0-3073. Reproduction, trans-
lation, publication, use and disposal, in whole or in
part, by or for the United States Government is permitted.

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America.

Available from:

National Technical Information Service
U. S. Department of Commerce
5285 Port Royal Road
Springfield, Virginia 22151

Price: Printed Copy \$ * ; Microfische \$3.50

<u>*PAGES</u>	<u>NTIS Selling Price</u>
1-25	\$5.00
26-50	\$6.50
51-75	\$8.00
76-100	\$9.50
101-125	\$11.00
126-150	\$12.50
151-175	\$14.00
176-200	\$15.50
201-225	\$17.00
226-250	\$18.50
251-275	\$20.00
276-300	\$21.50
301-325	\$23.00
326-350	\$24.50
351-375	\$26.00
376-400	\$27.50
401-425	\$29.00
426-450	\$30.50
451-475	\$32.00
476-500	\$33.50
500-525	\$35.00
526-550	\$36.50
551-575	\$38.00
576-600	\$39.50

For documents over 600 pages, add \$1.50 for each additional 25 page increment.

Hamiltonian Field Description of
Two-Dimensional Vortex Fluids and
Guiding Center Plasmas

by

Philip J. Morrison

Plasma Physics Laboratory, Princeton University

Princeton, New Jersey 08544

Abstract

The equations that describe the motion of two-dimensional vortex fluids and guiding center plasmas are shown to possess underlying field Hamiltonian structure. A Poisson bracket which is given in terms of the vorticity, the physical although noncanonical dynamical variable, casts these equations into Heisenberg form. The Hamiltonian density is the kinetic energy density of the fluid. The well-known conserved quantities are seen to be in involution with respect to this Poisson bracket. Expanding the vorticity in terms of a Fourier-Dirac series transforms the field description given here into the usual canonical equations for discrete vortex motion. A Clebsch potential representation of the vorticity transforms the noncanonical field description into a canonical description.

I. Introduction

This paper is concerned with the Hamiltonian field formulation of the equations which describe the advection of vorticity in a two-dimensional fluid. These equations have received a great deal of attention in the past thirty years and are believed to model the large scale motions which occur in atmospheres and oceans. They have also arisen in the study of plasma transport perpendicular to a uniform magnetic field, the so-called guiding center plasma.^{1,2} (For recent reviews see Refs. 3 and 4.)

It has been known for some time that a system of discrete vortex (or charge) filaments possesses a Hamiltonian description.⁵ The equations of motion are

$$k_i \frac{dx_i}{dt} = \frac{\partial H}{\partial y_i} \quad k_i \frac{dy_i}{dt} = - \frac{\partial H}{\partial x_i} \quad (1)$$

where k_i is the circulation of the i^{th} vortex which has coordinates x_i and y_i . The Hamiltonian, H , is the interaction energy and for an unbounded fluid has the form

$$H = \frac{-1}{2\pi} \sum_{i>j} k_i k_j \ln R_{ij},$$

where $R_{ij}^2 = (x_i - x_j)^2 + (y_i - y_j)^2$. The variables x_i and y_i are canonically conjugate. The formulation we describe here is a field formulation which possesses this underlying discrete dynamics.

In Sec. II we briefly review some aspects of finite degree of freedom Hamiltonian dynamics. The emphasis here is placed on the Lie algebraic properties of the Poisson bracket. This is used as a framework in which to

explain the "constructive" approach to Hamiltonian dynamics. Such an approach frees one from the prejudice that a system need be in canonical variables to be Hamiltonian. This section is then concluded by the extension of these notions to infinite degree of freedom or Hamiltonian density systems. In Sec. III we present a Poisson bracket that renders the vortex equations into Heisenberg form. This formulation is novel in that it is noncanonical. In the remainder of this section we discuss involutivity of the well-known constants of motion for this system, Fourier space representation and truncation. In Sec. IV we expand the vorticity in a Fourier-Dirac series which, upon substitution into the Poisson bracket of Sec. III, yields the canonical discrete vortex description of the Introduction. Following this we introduce Clebsch potentials which also bring the Poisson bracket into canonical form. Finally, we obtain a spectral description where complex conjugate pairs are canonically conjugate. A quartic interaction Hamiltonian is obtained.

II. Constructive Hamiltonian Dynamics

The standard approach⁶ to a Hamiltonian description is via a Lagrangian description. One constructs a Lagrangian on physical bases and through the Legendre transformation (assuming convexity) obtains the Hamiltonian and the following $2N$ (where N is the number of degrees of freedom) first order ordinary differential equations:

$$\dot{q}_k = [q_k, H] ; \quad \dot{p}_k = [p_k, H] \quad k = 1, 2, \dots, N \quad . \quad (2)$$

Here the Poisson bracket has the form

$$[f, g] = \sum_{k=1}^N \left(\frac{\partial f}{\partial q_k} \frac{\partial g}{\partial p_k} - \frac{\partial f}{\partial p_k} \frac{\partial g}{\partial q_k} \right) = \frac{\partial f}{\partial Z^i} J^{ij} \frac{\partial g}{\partial Z^j} \quad (3)$$

The last equality of Eq. (4) follows from the substitutions,

$$Z^i = \begin{cases} q_k & \text{for } i = k = 1, 2, \dots, N \\ p_k & \text{for } i = N + k = (N + 1) \dots 2N \end{cases}$$

and

$$(J^{ij}) = \begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix} \quad (4)$$

where I_N is the $N \times N$ unit matrix. We assume the repeated index summation convention here and henceforth. The quantity J^{ij} is known as the Poisson tensor or the cosymplectic form. It is not difficult to show that it transforms as a contravariant tensor under a change of coordinates. Those transformations which preserve its form, and hence the form of the Eqs. (2), the equations of motion, are canonical.

The constructive approach differs from the above in that one is not concerned with any underlying action principle nor with (initially at least) the necessity of canonical variables. The emphasis is placed on the algebraic properties of the Poisson bracket. A system need not have the canonical form of Eqs. (2) with Eq. (3) to be Hamiltonian. To make the idea more precise we introduce a few mathematical concepts. The quantities on which the Poisson bracket acts are differentiable functions defined on phase space. The collection of all such functions is a vector space (call it Ω) under addition and scalar multiplication. The Poisson bracket is a bilinear function which maps $\Omega \times \Omega$ to Ω . Also note that the Poisson bracket possesses the following

two important properties: (i) $[f,g] = -[g,f]$ for every $f,g \in \Omega$ and (ii) the Jacobi identity, i.e., $[f,[g,h]] + [g,[h,f]] + [h,[f,g]] = 0$ for every $f,g,h \in \Omega$. A vector space together with such a bracket defines a Lie algebra. Property (i) requires that the Poisson tensor be antisymmetric and property (ii) requires the following:

$$s^{ijk} \equiv J^{i\ell} \frac{\partial}{\partial Z^\ell} J^{jk} + J^{j\ell} \frac{\partial}{\partial Z^\ell} J^{ki} + J^{k\ell} \frac{\partial}{\partial Z^\ell} J^{ij} = 0 \quad . \quad (5)$$

One can show that s^{ijk} transforms contravariantly; hence if it vanishes identically in one coordinate frame it does so in all. Similarly antisymmetry is coordinate independent. The covariance of properties (i) and (ii) suggests the converse outlook: if a system of equations possesses the form

$$\dot{z}^i = \tilde{J}^{ij} \frac{\partial H}{\partial z^j} \quad i,j = 1,2,\dots,2N \quad (6)$$

where \tilde{J}^{ij} is antisymmetric and fulfills the Jacobi requirement, but is not of the form of Eq. (4), then it is Hamiltonian. This outlook is justified by a theorem due to Darboux (1882) which states that assuming $\det(J^{ij}) \neq 0$ (locally) canonical coordinates can be constructed. (The proof of this theorem may be found in Refs. 7, 8, and 9.) Hence in order for a system to be Hamiltonian it is only necessary for it be representable in Heisenberg form with a Poisson bracket that makes Ω into a Lie algebra. The constructive approach simply amounts to constructing Poisson brackets with the appropriate properties.

The rigorous generalization of the above ideas to infinite dimensional systems requires the language of functional analysis and the differential geometry of infinite dimensional manifolds. (See Ref. 7, Ch. V and Refs. 10 -

15.) This of course is not our purpose here; rather we simply parallel the above. The Poisson bracket for a set of field equations usually has the following form:⁶

$$[\hat{F}, \hat{G}] = \sum_{k=1}^N \int \left(\frac{\delta \hat{F}}{\delta \eta_k} \frac{\delta \hat{G}}{\delta \pi_k} - \frac{\delta \hat{G}}{\delta \eta_k} \frac{\delta \hat{F}}{\delta \pi_k} \right) d\tau \quad (7)$$

where the integration is taken over a fixed volume. The quantities on which the bracket acts are now functionals, such as the integral of the Hamiltonian density [e.g., Eq.(13)]. The functional derivative is defined by

$$\left. \frac{d\hat{F}}{d\varepsilon} (\eta_k + \varepsilon w) \right|_{\varepsilon=0} = \int \frac{\delta \hat{F}}{\delta \eta_k} w d\tau \equiv \langle \frac{\delta \hat{F}}{\delta \eta_k} | w \rangle$$

where the bra-ket notation is used to indicate the inner product $\langle f | g \rangle = \int f g d\tau$. In terms of this notation Eq. (7) becomes

$$[\hat{F}, \hat{G}] = \left\langle \frac{\delta \hat{F}}{\delta u^i} \mid O^{ij} \frac{\delta \hat{G}}{\delta u^j} \right\rangle \quad (8)$$

where the $2N$ quantities η_k and π_k are as previously the $2N$ indexed quantities u^i . The canonical cosymplectic density has the form

$$(O^{ij}) = \begin{pmatrix} 0 & I_N \\ -I_N & 0 \end{pmatrix} .$$

In noncanonical variables the quantity (O^{ij}) may depend upon the variables u^i , and further it may contain derivatives with respect to the independent variables. In general antisymmetry of Eq. (8) requires that the (O^{ij}) be an anti-self-adjoint operator. The Jacobi identity places further restrictions, analogous to Eq. (5), on this quantity. We defer a discussion of this to the

Appendix where the Jacobi identity for the bracket we present [Eq. (15)] is proved. The extension of the Darboux theorem to infinite dimensions has been proved by J. Marsden.¹⁴ For a discussion pertinent here see Ref. 15.

III. Noncanonical Poisson Bracket

The equations under consideration are the following:

$$\omega_t = - \underline{v} \cdot \nabla \omega \quad (9)$$

$$\nabla \cdot \underline{v} = 0 \quad (10)$$

Here we use the usual Euclidian coordinate system with uniformity in the z direction (which has unit vector \hat{z}). The quantity $\omega(\underline{x}, t) = \hat{z} \cdot \nabla \times \underline{v}(\underline{x}, t)$, where $\underline{x} \equiv (x, y)$, is the vorticity and \underline{v} is the flow velocity such that $\underline{v} \cdot \hat{z} = 0$. (For the guiding center plasma ω corresponds to the charge density and \underline{v} to the $\underline{E} \times \underline{B}$ drift velocity.) For an unbounded fluid \underline{v} can be eliminated from Eq. (9) by¹⁶

$$\underline{v} = \int \omega(\underline{x}') \underline{M}(\underline{x}|\underline{x}') d\tau' \quad (11)$$

where we display only the arguments necessary to avoid confusion. Here $\underline{M} = \hat{z} \times \nabla K(\underline{x}|\underline{x}')$ and $K(\underline{x}|\underline{x}')$ is the Green function for Laplace's equation in two dimensions,

$$K(\underline{x}|\underline{x}') = \frac{1}{2\pi} \ln \sqrt{(x-x')^2 + (y-y')^2} \quad .$$

The integration in Eq. (11) is over the entire x-y plane; $d\tau \equiv dx dy$. In this form Eq. (10) is satisfied manifestly. Equation (9) becomes

$$\omega_t = - \int \omega(\underline{x}') \underline{M}(\underline{x}|\underline{x}') d\tau' \cdot \nabla \omega(\underline{x}) \quad . \quad (12)$$

Equations (9) and (10) are known to possess conserved densities; that is, quantities which satisfy an equation of the form $\rho_t + \nabla \cdot \underline{J} = 0$, consistent with Eq. (12). Clearly any function of ω is conserved. In addition, the kinetic energy is conserved which is the natural choice for the Hamiltonian. With the density (mass) set to unity we have

$$\begin{aligned} \hat{H}\{\omega\} &= \int \frac{v^2}{2} d\tau = \frac{1}{2} \int \underline{M}(\underline{x}|\underline{x}') \cdot \underline{M}(\underline{x}|\underline{x}'') \omega(\underline{x}') \omega(\underline{x}'') d\tau d\tau' d\tau'' \\ &= - \frac{1}{2} \int K(\underline{x}|\underline{x}') \omega(\underline{x}') \omega(\underline{x}) d\tau d\tau' \quad . \end{aligned} \quad (13)$$

The functional derivative of Eq. (13) is the following:

$$\frac{\delta \hat{H}}{\delta \omega} = - \int K(\underline{x}|\underline{x}') \omega(\underline{x}') d\tau' \quad . \quad (14)$$

We introduce the Poisson bracket¹⁷

$$[\hat{F}, \hat{G}] = \int \omega(\underline{x}) \left\{ \frac{\delta \hat{F}}{\delta \omega}, \frac{\delta \hat{G}}{\delta \omega} \right\} d\tau \quad , \quad (15)$$

where $\{f, g\} = \frac{\partial f}{\partial x} \frac{\partial g}{\partial y} - \frac{\partial g}{\partial y} \frac{\partial f}{\partial x}$. One observes that the discrete vortex Poisson bracket is nestled inside the field Poisson bracket. In Sec. IV we will see how to regain the discrete bracket from this field bracket. It is not difficult to show from Eqs. (14) and (15) that

$$\omega_t = [\omega, \hat{H}] = - \int \omega \underline{M} \, d\tau' \cdot \nabla \omega \quad .$$

Clearly this bracket is antisymmetric by virtue of the antisymmetry of the discrete bracket. We prove the Jacobi identity in the Appendix.

We note by examination of Eq. (15) that any two functionals of ω are in involution; that is, if $\hat{F}_i\{\omega\} = \int F_i(\omega) d\tau$ (for $i=1,2$) are two such functionals where the F_i are arbitrary functions of ω , then

$$[\hat{F}_1, \hat{F}_2] = 0 \quad .$$

Also, substitution of any such \hat{F}_i and \hat{H} [Eq. (13)] into Eq. (15) and integration by parts yields

$$[\hat{F}_i, \hat{H}] = 0 \quad .$$

In particular, we see (when $F_i = \omega^2$) that the enstrophy commutes with the Hamiltonian.

The close relationship between this functional Hamiltonian formulation and the conventional formulation of Sec. II is seen by Fourier expanding the vorticity in a unit box with periodic boundary conditions,

$$\omega = \sum_{\underline{k}} \omega_{\underline{k}} e^{i\underline{k} \cdot \underline{x}} \quad , \quad (16)$$

where $\underline{k} = (k_1, k_2)$. The reality of ω implies $\omega_{\underline{k}}^* = \omega_{-\underline{k}}$. If we suppose for the moment that $\omega(\underline{x})$ depends upon some additional independent variable μ , then we have the following for some functional \hat{F} :

$$\frac{\partial \hat{F}}{\partial \mu} = \int \frac{\delta \hat{F}}{\delta \omega} \frac{\partial \omega}{\partial \mu} dx dy \quad . \quad (17)$$

From this we see for $\mu = \omega_{\underline{k}}$ upon Fourier inversion that

$$\frac{\delta \hat{F}}{\delta \omega} = \frac{1}{(2\pi)^2} \sum_{\underline{k}} \frac{\partial \hat{F}}{\partial \omega_{\underline{k}}} e^{-i\underline{k} \cdot \underline{x}} \quad , \quad (18)$$

where the \hat{F} on the left hand side is treated as a functional of ω while the \hat{F} of the right hand side is to be regarded as a function of the variables $\omega_{\underline{k}}$. Substituting Eqs. (16) and (18) into Eq. (15) yields,

$$[\hat{F}, \hat{G}] = \sum_{\underline{k}, \underline{l}} \frac{\omega_{\underline{k}+\underline{l}}}{(2\pi)^2} \hat{z} \cdot (\underline{l} \times \underline{k}) \frac{\partial \hat{F}}{\partial \omega_{\underline{k}}} \frac{\partial \hat{G}}{\partial \omega_{\underline{l}}} \quad .$$

The Hamiltonian becomes

$$\hat{H} = 2\pi^2 \sum_{\underline{l}} \frac{|\omega_{\underline{l}}|^2}{l^2} \quad ,$$

and the equations of motion are

$$\dot{\omega}_{\underline{k}} = \sum_{\underline{l}} \frac{\hat{z} \cdot (\underline{k} \times \underline{l})}{l^2} \omega_{\underline{l}} \omega_{\underline{k}-\underline{l}} = \sum_{\underline{l}} J_{\underline{k}, \underline{l}} \frac{\partial \hat{H}}{\partial \omega_{\underline{l}}} \quad , \quad (19)$$

where $J_{\underline{k}, \underline{l}}$, the cosymplectic form, is

$$J_{\underline{k}, \underline{l}} = \frac{\hat{z} \cdot (\underline{l} \times \underline{k})}{(2\pi)^2} \omega_{\underline{k}+\underline{l}} \quad . \quad (20)$$

Clearly, Eq. (19) is of the form of the finite degree of freedom equations, Eqs. (6), of Sec. II except here the sum ranges to infinity. The form Eq.

(20) is obviously antisymmetric and it is not difficult to verify Eq. (5).

At first, one might think that a truncation of the $J_{\underline{k}, \underline{\ell}}$ would yield a finite Hamiltonian system which to some accuracy would mimic the original. Unfortunately, the process of truncation destroys the Jacobi identity. One must seek a change of variables which allows truncation. Canonical variables are suited for this purpose and in the next section we discuss this.

IV. Canonical Descriptions

As was noted in Sec. III, the Poisson bracket for the discrete vortex picture is embedded in that for the field. To see the connection between the two, we expand the vorticity (distributed vorticity) as follows:

$$\omega(\underline{x}) = \sum_i k_i \delta(\underline{x} - \underline{x}_i) \quad , \quad (21)$$

where $\delta(\underline{x})$ is the Dirac delta function, the k_i are constants and ω obtains its t dependence through the \underline{x}_i . Then using Eq. (17) we obtain the identity

$$\left. \frac{\partial \hat{F}}{\partial \underline{x}_i} = k_i \frac{\partial}{\partial \underline{x}} \frac{\delta \hat{F}}{\delta \omega} \right|_{\underline{x} = (\underline{x}_i, y_i)} \quad (22)$$

where the functional \hat{F} on the left hand side is now to be regarded as a function of the variables x_i and y_i . Similarly we obtain the relation for $\partial \hat{F} / \partial y_i$. Substituting Eqs. (21) and (22) into Eq. (15) yields

$$[\hat{F}, \hat{G}] = \sum_j \frac{1}{k_j} \left(\frac{\partial \hat{F}}{\partial x_j} \frac{\partial \hat{G}}{\partial y_j} - \frac{\partial \hat{F}}{\partial y_j} \frac{\partial \hat{G}}{\partial x_j} \right) \quad . \quad (23)$$

Further, if we substitute Eq. (21) into the Hamiltonian, Eq. (13), we obtain

$$\hat{H} = \frac{-1}{4\pi} \sum_{i,j} k_i k_j \ln R_{ij} \quad .$$

Since this is singular along the diagonal $i=j$, we remove the self-energy of each vortex and obtain the usual result

$$\hat{H} = \frac{-1}{2\pi} \sum_{i>j} k_i k_j \ln R_{ij} \quad . \quad (24)$$

Equations (23) and (24) reproduce the Eqs. (1). Hence, we see that expansion of ω in a Fourier-Dirac series is a particular way of discretizing, a way which allows truncation without destroying the Hamiltonian structure. We now discuss another approach.

The cosymplectic form, Eq. (20), suggests by its linearity in $\omega_{\underline{k}+\underline{\ell}}$ that a quadratic change of variables (i.e., $\omega \sim \phi^2$, where ϕ is the new variable) is needed in order to achieve canonical form. Such a transformation [given by Eq. (31)] removes the nonlinearity present in the Poisson bracket and places it in the Hamiltonian [Eq. (30)]. Enroute to arriving at this result we introduce a Clebsch potential representation of the vorticity,²¹

$$\omega = \frac{\partial \psi}{\partial x} \frac{\partial \chi}{\partial y} - \frac{\partial \psi}{\partial y} \frac{\partial \chi}{\partial x} \quad . \quad (25)$$

This substitution transforms the Poisson bracket, Eq. (15), into canonical form. Clearly, Eq. (25) is not uniquely invertable. We have the local gauge condition that any function $\bar{\psi}$, such that $\bar{\psi}_x \chi_y - \bar{\psi}_y \chi_x = 0$, can be added to ψ (and likewise for χ).

The chain rule for functional differentiation yields

$$\frac{\hat{\delta F}}{\delta \psi} = \nabla \cdot \left(\frac{\hat{\delta F}}{\delta \omega} \hat{z} \times \nabla \chi \right)$$

$$\frac{\hat{\delta F}}{\delta \chi} = - \nabla \cdot \left(\frac{\hat{\delta F}}{\delta \omega} \hat{z} \times \nabla \psi \right) \quad , \quad (26)$$

where on the left \hat{F} is now regarded as a functional of ψ and χ . The canonical Poisson bracket for χ and ψ is

$$[\hat{F}, \hat{G}] = \int \left(\frac{\hat{\delta F}}{\delta \psi} \frac{\hat{\delta G}}{\delta \chi} - \frac{\hat{\delta G}}{\delta \psi} \frac{\hat{\delta F}}{\delta \chi} \right) d\tau \quad ,$$

which upon substitution of Eqs. (26) yields the bracket Eq. (15). Clearly ψ and χ satisfy

$$\dot{\psi} = \frac{\hat{\delta H}}{\delta \chi} \quad ; \quad \dot{\chi} = - \frac{\hat{\delta H}}{\delta \psi} \quad . \quad (27)$$

Upon Fourier transformation Eqs. (27) become

$$\dot{\psi}_{\underline{k}} = \frac{1}{(2\pi)^2} \frac{\partial \hat{H}}{\partial \chi_{-\underline{k}}} \quad ; \quad \dot{\chi}_{\underline{k}} = - \frac{1}{(2\pi)^2} \frac{\partial \hat{H}}{\partial \psi_{-\underline{k}}} \quad . \quad (28)$$

We now introduce the field variable $\phi_{\underline{k}}$ as follows:

$$\psi_{\underline{k}} = \frac{1}{2\pi} \left(\frac{\phi_{\underline{k}} + \phi_{-\underline{k}}^*}{\sqrt{2}} \right) \quad ; \quad \chi_{\underline{k}} = \frac{-i}{2\pi} \left(\frac{\phi_{\underline{k}} - \phi_{-\underline{k}}^*}{\sqrt{2}} \right) \quad .$$

(This form maintains the reality condition for $\psi_{\underline{k}}$ and $\chi_{\underline{k}}$.) In terms of these variables Eqs. (28) become

$$i\dot{\phi}_{\underline{k}} = \frac{\hat{\delta H}}{\partial \phi_{\underline{k}}^*} \quad ; \quad i\dot{\phi}_{\underline{k}}^* = - \frac{\hat{\delta H}}{\partial \phi_{\underline{k}}} \quad , \quad (29)$$

and the Hamiltonian has the form

$$\hat{H} = \sum_{\underline{\ell} + \underline{m} = \underline{s} + \underline{t}} S_{\underline{\ell}, \underline{m}, \underline{s}, \underline{t}} \phi_{\underline{\ell}}^* \phi_{\underline{m}}^* \phi_{\underline{s}} \phi_{\underline{t}} \quad , \quad (30)$$

where the matrix elements $S_{\underline{\ell}, \underline{m}, \underline{s}, \underline{t}}$ are

$$S_{\underline{\ell}, \underline{m}, \underline{s}, \underline{t}} = \frac{1}{16\pi^2} \left[\frac{\hat{z} \cdot (\underline{t} \times \underline{\ell}) \hat{z} \cdot (\underline{m} \times \underline{s})}{|\underline{\ell} - \underline{t}| |\underline{m} - \underline{s}|} + \frac{\hat{z} \cdot (\underline{s} \times \underline{\ell}) \hat{z} \cdot (\underline{m} \times \underline{t})}{|\underline{\ell} - \underline{m}| |\underline{m} - \underline{t}|} \right] .$$

The quadratic transformation mentioned above is

$$\omega_{\underline{k}} = \sum_{\underline{t} = \underline{k} + \underline{\ell}} \frac{iz \cdot (\underline{t} \times \underline{\ell})}{(2\pi)^2} \phi_{\underline{t}} \phi_{\underline{\ell}}^* . \quad (31)$$

Hence, we see the connection between Clebsch potentials and our bracket. This transformation allows discretization and truncation while not destroying the Hamiltonian structure.

Acknowledgments

I would like to thank J. M. Greene for his continuing support and encouragement of this work. Also, I would like to thank H. Segur for stimulating my interest in the 2-D fluid equations. I am grateful to C. Oberman, J. B. Taylor, and G. Sandri for many discussions which contributed to this paper. I would like to thank J. E. Marsden for sending me an early copy of Ref. 15.

This work was supported by the U.S. Department of Energy Contract No. DE-ACO2-76-CHO3073.

Appendix

Here we generalize the method used by P. Lax²² for the Gardner bracket, to prove the Jacobi identity for Eq. (15). We suppose $\hat{F}\{u\}$ is a functional of the variable u . Recall the functional derivative is defined by

$$\frac{d}{d\varepsilon} \hat{F}\{u + \varepsilon\omega\} \Big|_{\varepsilon=0} = \left\langle \frac{\delta \hat{F}}{\delta u} \mid \omega \right\rangle .$$

We denote $\hat{G} = \left\langle \frac{\delta \hat{F}}{\delta u} \mid \omega \right\rangle$. \hat{G} can again be regarded as a functional of u . Performing a second variation we obtain

$$\frac{d}{d\eta} \hat{G}\{u + \eta z\} \Big|_{\eta=0} = \left\langle \frac{\delta^2 \hat{F}}{\delta u^2} z \mid \omega \right\rangle$$

where the symbol $\delta^2 \hat{F} / \delta u^2$ is used to denote an operator acting on z . By the equality of mixed partial derivatives this operator is self-adjoint,

$$\left\langle \frac{\delta^2 \hat{F}}{\delta u^2} z \mid \omega \right\rangle = \left\langle z \mid \frac{\delta^2 \hat{F}}{\delta u^2} \omega \right\rangle .$$

Let us now take the variation of a bracket $[\hat{F}, \hat{G}]$ defined by

$$[\hat{F}, \hat{G}] = \left\langle \frac{\delta \hat{F}}{\delta u} \mid O \frac{\delta \hat{G}}{\delta u} \right\rangle ,$$

where the operator O is anti-self-adjoint.

$$\begin{aligned} \frac{d}{d\varepsilon} [\hat{F}, \hat{G}]\{u + \varepsilon\omega\} \Big|_{\varepsilon=0} &= \left\langle \frac{\delta [\hat{F}, \hat{G}]}{\delta u} \mid \omega \right\rangle \\ &= \left\langle \frac{\delta^2 \hat{F}}{\delta u^2} \omega \mid O \frac{\delta \hat{G}}{\delta u} \right\rangle + \left\langle \frac{\delta \hat{F}}{\delta u} \mid O \frac{\delta^2 \hat{G}}{\delta u^2} \omega \right\rangle + \left\langle \frac{\delta \hat{F}}{\delta u} \mid \frac{\delta O}{\delta u} \omega \frac{\delta \hat{G}}{\delta u} \right\rangle . \end{aligned}$$

In this expression the first two terms are straight forward, the last comes from any dependence the operator O may have upon u ; i.e.,

$$\frac{d}{d\varepsilon} O(u + \varepsilon w) \Big|_{\varepsilon=0} \equiv \frac{\delta O}{\delta u} w .$$

Isolating w we obtain

$$\frac{\delta[\hat{F}, \hat{G}]}{\delta u} = \frac{\delta^2 \hat{F}}{\delta u^2} O \frac{\delta \hat{G}}{\delta u} - \frac{\delta^2 \hat{G}}{\delta u^2} O \frac{\delta \hat{F}}{\delta u} + T \left(\frac{\delta \hat{F}}{\delta u}, \frac{\delta \hat{G}}{\delta u} \right) \quad (A-1)$$

where the operator T comes from removing $\delta O_w / \delta u$ from w . T is antisymmetric in its arguments.

The Jacobi identity is

$$S = \left\langle \frac{\delta \hat{E}}{\delta u} \mid O \frac{\delta[\hat{F}, \hat{G}]}{\delta u} \right\rangle + \left\langle \frac{\delta \hat{F}}{\delta u} \mid O \frac{\delta[\hat{G}, \hat{E}]}{\delta u} \right\rangle + \left\langle \frac{\delta \hat{G}}{\delta u} \mid O \frac{\delta[\hat{E}, \hat{F}]}{\delta u} \right\rangle = 0 . \quad (A-2)$$

Inserting Eq. (A-1) into (A-2) and using the self-adjointness of $\frac{\delta^2 \hat{F}}{\delta u^2}$ and the anti-self-adjointness of O , we obtain

$$S = \left\langle \frac{\delta \hat{E}}{\delta u} \mid O T \left(\frac{\delta \hat{F}}{\delta u}, \frac{\delta \hat{G}}{\delta u} \right) \right\rangle + \left\langle \frac{\delta \hat{F}}{\delta u} \mid O T \left(\frac{\delta \hat{G}}{\delta u}, \frac{\delta \hat{E}}{\delta u} \right) \right\rangle + \left\langle \frac{\delta \hat{G}}{\delta u} \mid O T \left(\frac{\delta \hat{E}}{\delta u}, \frac{\delta \hat{F}}{\delta u} \right) \right\rangle = 0 . \quad (A-3)$$

This equation is the functional equivalent of Eq. (5).

Now consider the bracket, Eq. (15). We obtain

$$\frac{\delta[\hat{F}, \hat{G}]}{\delta \omega} = \left\{ \frac{\delta \hat{F}}{\delta \omega}, \frac{\delta \hat{G}}{\delta \omega} \right\} + \text{operator terms,}$$

where the first term is $T \left(\frac{\hat{\delta F}}{\delta \omega}, \frac{\hat{\delta G}}{\delta \omega} \right)$ and the remaining terms as shown above do not enter Eq. (A-3). Hence,

$$S = \int \omega \left\{ \frac{\hat{\delta E}}{\delta \omega}, \left\{ \frac{\hat{\delta F}}{\delta \omega}, \frac{\hat{\delta G}}{\delta \omega} \right\} \right\} dt + \text{cyc} \quad .$$

Clearly, S vanishes by virtue of the Jacobi identity for the discrete bracket.

References

1. J. B. Taylor and B. McNamara, *Phys. of Fluids* 14, 1492 (1971).
2. G. Vahala and D. Montgomery, *J. Plasma Physics* 6, 425 (1971).
3. P. Rhines, *Ann. Rev. Fluid Mech.* 11, 401 (1979).
4. P. H. Kraichnan and D. Montgomery, *Rep. Prog. Phys.* 43, 547 (1980).
5. In Ref. 4 they cite study on discrete vortices as early as 1858 by Helmholtz. More recent references are: C. C. Lin, On the motion of vortices in two dimensions, (University of Toronto Press, Toronto, 1943); L. Onsager, *Nuovo Cim. Suppl.* 6, 279 (1949).
6. H. Goldstein, Classical Mechanics, (Addison-Wesley, Reading, Mass., 1950). For more rigorous approaches see Refs. 8 and 9.
7. R. Abraham and J. E. Marsden, Foundations of Mechanics, (Benjamin-Cummings, London, 1978).
8. V. I. Arnold, Mathematical Methods of Classical Mechanics, (Springer-Verlag, New York, 1978).
9. R. Littlejohn, *J. Math. Phys.* 20, 2445 (1979).

10. P. R. Chernoff and J. E. Marsden, Properties of Infinite Dimensional Hamiltonian Systems, Lect. Notes in Math. 425 (Springer-Verlag, Berlin, 1974).

11. Y. I. Manin, J. Sov. Math. 11, 1 (1979).

12. I. M. Gel'fand and I. Y. Dorfman, Func. Anal. Appl. 13, 248 (1979).

13. B. Kupperschmidt, in Lect. Notes in Mathematics 755, 162; Edited by A. Dold and B. Eckmann, (Springer-Verlag, Berlin, 1980).

14. J. E. Marsden, Lect. Notes on Geometric Meths. in Math. Phys., SIAM (1981).

15. J. E. Marsden and A. Weinstein, to be published.

16. This formula is not unique. Without damage to the forthcoming Hamiltonian structure one may add to \underline{y} the cross product of \hat{z} with the gradient of an arbitrary harmonic function.

17. Clearly, this bracket is noncanonical. It appears that the first use of a noncanonical bracket was by C. S. Gardner (Ref. 18) where a bracket for the Korteweg-deVries equation was given. Recently brackets for an ideal fluid and MHD were obtained in Ref. 19. Brackets for the Maxwell-Vlasov and Poisson-Vlasov equations were obtained in Ref. 20. (See Ref. 15 for generalization.) We note that the Poisson bracket given here [Eq. (15)] is precisely that for the one-dimensional Vlasov-Poisson equations if one

replaces the vorticity by the distribution function and the phase space (x,y) by (x,v) , although the Hamiltonians are different. This is the subject of, P. J. Morrison, Princeton Plasma Phys. Lab Report #1788 (1981), to be published.

18. C. S. Gardner, J. Math. Phys. 12, 1548 (1971).

19. P. J. Morrison and J. M. Greene, Phys. Rev. Lett. 45, 790 (1980).

20. P. J. Morrison, Phys. Lett. 80A, 383 (1980).

21. A. Clebsch, Crelle, J. Reine Angew. Math. 56, 1 (1859). See H. Lamb. Hydrodynamics, sixth edition, (Cambridge University Press, Cambridge, 1932) sec. 167; and J. Serrin, in Handbuch der Physik, ed. by S. Flugge (Springer-Verlag, Berlin, 1959), Vol. 8 p. 169. Also, for three-dimensional compressible hydrodynamics see V. E. Zakharov, Sov. Phys. JETP 33, 927 (1971). For this case equations analogous to Eqs. (29) are given.

22. P. Lax, Comm. Pure Appl. Math. 28, 141 (1975).

ALL CATEGORIES

R. Askew, Auburn University, Alabama
 S. T. Wu, Univ. of Alabama
 Geophysical Institute, Univ. of Alaska
 G.L. Johnston, Sonoma State Univ, California
 H. H. Kuehl, Univ. of S. California
 Institute for Energy Studies, Stanford University
 H. D. Campbell, University of Florida
 N. L. Oleson, University of South Florida
 W. M. Stacey, Georgia Institute of Technology
 Benjamin Ma, Iowa State University
 Magne Kristiansen, Texas Tech. University
 W. L. Wiese, Nat'l Bureau of Standards, Wash., D.C.
 Australian National University, Canberra
 C.N. Watson-Munro, Univ. of Sydney, Australia
 F. Cap, Inst. for Theo. Physics, Austria
 Dr.M. Heindler, Institute for Theoretical Physics
 Technical University of Graz
 Ecole Royale Militaire, Bruxelles, Belgium
 D. Palumbo, C. European Comm. B-1049-Brussels
 P.H. Sakanaka, Instituto de Fisica, Campinas, Brazil
 M.P. Bachynski, MPB Tech., Ste. Anne de Bellevue,
 Quebec, Canada
 C. R. James, University of Alberta, Canada
 T.W. Johnston, INRS-Energie, Varennes, Quebec
 H. M. Skarsgard, Univ. of Saskatchewan, Canada
 Inst. of Physics, Academia Sinica, Peking,
 People's Republic of China
 Inst. of Plasma Physics, Hefei,
 Anhwei Province, People's Republic of China
 Library, Tsing Hua Univ., Peking, People's
 Republic of China
 Zhengwu Li, Southwestern Inst. of Phys., Leshan,
 Sichuan Province, People's Republic of China
 Librarian, Culham Laboratory, Abingdon, England (2)
 A.M. Dupas Library, C.E.N.-G, Grenoble, France
 Central Res. Inst. for Physics, Hungary
 S. R. Sharma, Univ. of Rajasthan, JAIPUR-4, India
 R. Shingal, Meerut College, India
 A.K. Sundaram, Phys. Res. Lab., India
 Biblioteca, Frascati, Italy
 Biblioteca, Milano, Italy
 G. Rostagni, Univ. Di Padova, Padova, Italy
 Preprint Library, Inst. de Fisica, Pisa, Italy
 Library, Plasma Physics Lab., Gokasho, Uji, Japan
 S. Mori, Japan Atomic Energy Res. Inst., Tokai-Mura
 Research Information Center, Nagoya Univ., Japan
 S. Shioda, Tokyo Inst. of Tech., Japan
 Inst. of Space & Aero. Sci., Univ. of Tokyo
 T. Uchida, Univ. of Tokyo, Japan
 H. Yamato, Toshiba R. & D. Center, Japan
 M. Yoshikawa, JAERI, Tokai Res. Est., Japan
 Dr. Tsuneo Nakakita, Toshiba Corporation,
 Kawasaki-Ku Kawasaki, 210 Japan
 N. Yajima, Kyushu Univ., Japan
 R. England, Univ. Nacional Auto-noma de Mexico
 B. S. Liley, Univ. of Waikato, New Zealand
 S. A. Moss, Saab Univas Norge, Norway
 J.A.C. Cabral, Univ. de Lisboa, Portugal
 O. Petrus, A.L.I. CUZA Univ., Romania
 J. de Villiers, Atomic Energy Bd., South Africa
 A. Maurech, Comisaria De La Energy y Recursos
 Minerales, Spain
 Library, Royal Institute of Technology, Sweden
 Cen. de Res. En Phys.Des Plasmas, Switzerland
 Librarian, Fom-Instituut Voor Plasma-Fysica,
 The Netherlands

Bibliothek, Stuttgart, West Germany
 R.D. Buhler, Univ. of Stuttgart,
 West Germany
 Max-Planck-Inst. fur Plasmaphysik,
 W. Germany
 Nucl. Res. Estab., Julich, West Germany
 K. Schindler, Inst. Fur Theo. Physik,
 W. Germany

EXPERIMENTAL
THEORETICAL

M. H. Brennan, Flinders Univ. Australia
 H. Barnard, Univ. of British Columbia, Canada
 S. Screenivasan, Univ. of Calgary, Canada
 J. Radet, C.E.N.-B.P., Font enay-aux-Roses,
 France
 Prof. Schatzman, Observatoire de Nice,
 France
 S. C. Sharma, Univ. of Cape Coast, Ghana
 R. N. Aiyer, Laser Section, India
 B. Buti, Physical Res. Lab., India
 L. K. Chavda, S. Gujarat Univ., India
 I.M. Las Das, Banaras Hindu Univ., India
 S. Cuperman, Tel Aviv Univ., Israel
 E. Greenspan, Nuc. Res. Center, Israel
 P. Rosenau, Israel Inst. of Tech., Israel
 Int'l. Center for Theo. Physics, Trieste, Italy
 I. Kawakami, Nihon University, Japan
 T. Nakayama, Ritsumeikan Univ., Japan
 S. Nagao, Tohoku Univ., Japan
 J.I. Sakai, Toyama Univ., Japan
 S. Tjotta, Univ. I Bergen, Norway
 M.A. Hellberg, Univ. of Natal, South Africa
 H. Wilhelmson, Chalmers Univ. of Tech.,
 Sweden
 Astro. Inst., Sonnenborgh Obs.,
 The Netherlands
 T. J. Boyd, Univ. College of North Wales
 K. Hubner, Univ. Heidelberg, W.Germany
 H. J. Kaeppler, Univ. of Stuttgart,
 West Germany
 K. H. Spatschek, Univ. Essen, West Germany

EXPERIMENTAL
ENGINEERING

B. Grek, Univ. du Quebec, Canada
 P. Lukac, Komenskeho Univ., Czechoslovakia
 G. Horikoshi, Nat'l Lab for High Energy Physics,
 Tsukuba-Gun, Japan

EXPERIMENTAL

F. J. Paoloni, Univ. of Wollongong, Australia
 J. Kistemaker, Fom Inst. for Atomic
 & Molec. Physics, The Netherlands

THEORETICAL

F. Verheest, Inst. Vor Theo. Mech., Belgium
 J. Teichmann, Univ. of Montreal, Canada
 T. Kahan, Univ. Paris VII, France
 R. K. Chhajlani, India
 S. K. Trehan, Panjab Univ., India
 T. Namikawa, Osaka City Univ., Japan
 H. Narumi, Univ. of Hiroshima, Japan
 Korea Atomic Energy Res. Inst., Korea
 E. T. Karlson, Uppsala Univ., Sweden
 L. Stenflo, Univ. of UMEA, Sweden
 J. R. Saraf, New Univ., United Kingdom